

KİMYA TARIM ÜR. PAZ. SAN. ve TİC. LTD. ŞTİ. ●

nevagri



Product Catalog



The company was established in Antalya with the aim of producing and marketing food for plants. Our company produces chemical fertilizers, organic fertilizers and fertilizers in accordance with the needs of the agricultural sector.

We also produce organomineral fertilizers.

Following the innovations in the agricultural sector: our company attaches great importance to research and development.

By exploring innovations in the agricultural sector, we are working to improve our formulations. The source of our strength, which allows us to make steady progress, is our corporate quality policy.

For more information and our other products, please visit our website.

ORGANIC FERTILIZER LIQUID ORGANIC FERTILIZER CONTAINING AMINO ACIDS OF ANIMAL ORIGIN

LİSABLAN

GUARANTEED CONTENT	%W/W
Total organic matter	%60
Organic Nitrogen	%9
Organic Carbon	%28
Free Amino Acids	%11
pH	5,5 - 7,5

RECOMMENDATIONS

- Read the label carefully.
- Keep away from children, pets and food.
- Never exceed the appropriate dose and use only when necessary.
- Please use a mask, gloves and glasses during fertilization.
- Avoid contact with eyes, face and skin. In case of contact, rinse immediately with plenty of water.
- Do not use the used empty packaging for other purposes, destroy it.
- It is recommended to use it after analyzing the soil and/or leaves.

ATTENTION:

DO NOT USE IN SOIL ACCESSIBLE TO FARM ANIMALS.

DO NOT GRAZE OR LEAVE ANIMALS FOR AT LEAST 21 DAYS AFTER APPLICATION.

MOWN GRASS SHOULD NOT BE USED IN ANIMAL FEEDING.

WHAT SHOULD BE CONSIDERED WHEN USING AND STORING:

İSABON can be mixed with all fertilizers, chelates and most plant protection products. It should not be mixed with more than two other products. Mixing with mineral oils and nitro derivatives such as DNOC should be avoided. You can safely use a tank mixture with copper-containing fungicides on olives, cereals and onions. When using tank mixtures with sulfur and copper compounds, preliminary tests should be carried out on the target culture and monitor the reactions that may occur within a few days after application. This should be done 3-4 days after the application of copper compounds.

USE RECOMMENDATIONS

CROP	Application	FOLIAR APPLICATION	CROP	Application	FOLIAR APPLICATION
PISTACHIO	Before flowering, during the period of petal fall and fruiting	250 ml/100 L of water	CELERY	Apply every 20 days, starting from 10-15 days after planting	250 ml/ha
PEAR	Before flowering, when the petals fall off, during the fruiting period and when the fruits reach 6-7 cm in diameter	200-300 ml/100 L of water	CHESTNUT	Before flowering, during the period of petal fall and fruit setting	250 ml/100 L of water
BARLEY	From the 4-5 leaf phase to the end of tillering and 2 applications after tillering	300-500 ml/ha	CHERRY	Before flowering, when the petals fall off, during the fruiting period, when the fruits reach 6-7 cm in diameter	200-250 ml/100 L of water
SUNFLOWER	At the stage of 3-4 leaves and before flowering, 2 applications	300-500 ml/ha	KIWI	Vegetative development before flowering - When the petals fall off, during fruiting and when the fruits reach 6-7 cm in diameter	150-200 ml/100 L of water 250-300 ml/100 L of water
QUINCE	Before flowering, when the petals fall off, during the fruiting period and when the fruits reach 6-7 cm in diameter	200-300 ml/100 L of water	RAPESEED	During the formation of lateral branches and trunk elongation - 2 treatments	200 ml/ha
ALMOND	Before flowering, during petal fall and during fruiting	250 ml/100 L of water	THE CABBAGE FAMILY	After planting and two deposits with an interval of 20 days	250-300 ml/ha
THE VINEYARD	Before flowering, when the petals fall off and during the color change of the fruits	250-300 ml/100 L of water	LEMON	Before vegetative development and flowering - when the petals fall off, during the fruiting period and when the fruits reach 6-7 cm in diameter	200-300 ml/100 L of water
PEAS	Before flowering, 3 applications with an interval of 10-15 days After flowering	200-300 ml/100 L of water	MANDARIN	Before vegetative development and flowering - during the fall of the petals, during the fruiting period and when the fruits reach 6-7 cm in diameter	200-300 ml/100 L of water
PEPPER (field)	Periods before flowering, fruiting, and fruit ripening	250-300 ml/ha	LETTUCE (FIELD)	Application every 20 days 10-15 days after shaking	250 ml/ha
PEPPER (greenhouse)	Periods before flowering, fruiting, and fruit ripening	250-300 ml/100 L of water	LETTUCE (GREENHOUSE)	Application every 20 days 10-15 days after shaking	200 ml/100 L of water
WHEAT	From the 4-5 leaf phase to the end of tillering and after tillering - 2 applications	300-500 ml/ha	PUMPKIN	When the roots become the size of a pencil, 3-4 applications every 2-3 weeks	250-300 ml/ha
WALNUT	Before flowering, during the period of petal fall and fruiting	250 ml/100 L of water	LENTIL	25 days after sowing, 3 applications with an interval of 15 days	250-300 ml/ha
RYE	From the 4-5 leaf phase to the end of tillering and after tillering - 2 applications	300-500 ml/ha	CORN	25 days after sowing, 3 applications with an interval of 15 days	250-300 ml/ha
PODDY	2 applications during the tillering and flowering period.	1 application: 400-500 ml/ha 2 application: 500-600 ml/ha	BANANA	2-4 applications with an interval of 12-14 days during critical periods such as flowering, falling and fruit formation, discoloration, frost (use a lower dose on young plants aged 1-2 years)	250-350 ml/ha
STRAWBERRIES (field)	20-30 days before flowering, 3-5 days before fruit setting, every 20 days until the end of the production season	200-250 ml/ha	POMEGRANATE	Before flowering, when the petals fall off, during the fruiting period, when the fruits reach 6-7 cm in diameter	200-250 ml/100 L of water
STRAWBERRY (GREENHOUSE)	20-30 days before flowering at the beginning of vegetative development before hibernation, 3-5 days before fruit setting, every 20 days until the end of the production season	200-250 ml/100 L of water	NECTARINE	Before flowering, when the petals fall off, during the fruiting period, when the fruits reach 6-7 cm in diameter	200-250 ml/100 L of water
TOMATO (field)	Periods before flowering, fruiting, and fruit ripening	250-300 ml/ha	CHICKPEAS	25 days after sowing, 3 applications with an interval of 15 days, 3 applications before and 3 applications after flowering in the presence of 3-4 leaves	250-300 ml/ha
TOMATO (GREENHOUSE)	Periods before flowering, fruiting, and fruit ripening	250-300 ml/100 L of water	COTTON	When the plants reach 15 cm, when the tubers become the size of a handball, and after 15 days.	200-300 ml/ha
APPLE	Before flowering, when the petals fall off, during the fruiting period and when the fruits reach 6-7 cm in diameter	200-300 ml/100 L of water	POTATO	Before flowering, during fruiting, during fruit ripening, after the beginning of tuber formation	200-300 ml/ha
ARTICHOKE	3 days after planting and 4 applications every 15-20 days	200-300 ml/100 L of water	EGGPLANT (FIELD)	Fruiting before flowering and during fruit ripening	250-300 ml/ha
ERIC	Before flowering, when the petals fall off, during the fruiting period and when the fruits reach 6-7 cm in diameter	200-250 ml/100 L of water	EGGPLANT (GREENHOUSE)	Apply every 20 days, starting 10-15 days after planting	250-300 ml/100 L of water
BEANS	Before flowering, 3 applications with an interval of 10-15 days After flowering	300 ml/100 L of water	LEEK	Before vegetative development and flowering - during the fruiting period, when the petals fall off and when the fruits reach 6-7 cm in diameter	200 ml/ha
HAZELNUT	Before flowering, when the petals fall off and during fruiting	250 ml/100 L of water	ORANGE	A tank mixture with copper compounds every 20 days after the beginning of tuber formation	200-300 ml/100 L of water
GRAPEFRUIT	Vegetative development and flowering, when the petals fall off, during fruiting and when the fruits reach 6-7 cm in diameter	200-300 ml/100 L of water	GARLIC	A tank mixture with copper compounds every 20 days after the beginning of tuber formation	200-300 ml/100 L of water
CARROT	When the roots reach the size of a pencil, 3-4 applications every 2-3 weeks	300-400 ml/ha	ONION	A tank mixture with copper compounds every 20 days after the beginning of tuber formation	200-300 ml/100 L of water
CUCUMBER (FIELD)	Periods before flowering, fruiting, and fruit ripening	300 ml/ha	SOY	25 days after sowing, 3 applications with an interval of 15 days	250-300 ml/ha
CUCUMBER (GREENHOUSE)	Before flowering, fruit laying, fruit ripening periods	100-250 ml/100 L of water	PEACH	Before flowering, when the petals fall off, during the fruiting period, when the fruits reach 6-7 cm in diameter	200-250 ml/100 L of water
SPINACH	Application every 20 days, starting from 15-20 days after sowing	250 ml/ha	SUGAR BEET	One application every 20 days, starting from 15-20 days after sowing	250 ml/ha
PUMPKIN (FIELD)	4-5 leaves before flowering, fruit laying, fruit ripening periods	300 ml/ha	TOBACCO	One application every 20 days, starting 20 days after landing	200-250 ml/100 L of water
PUMPKIN (GREENHOUSE)	4-5 leaves before flowering, fruit setting, fruit ripening periods	100-250 ml/100 L of water	CHERRY	During the fruiting period, when the petals are shed before flowering, when the fruits reach 6-7 cm in diameter	200-250 ml/100 L of water
WATERMELON (FIELD)	4-5 leaves before flowering, fruit setting, fruit ripening periods	300 ml/ha	PEANUT	25 days after sowing, 3 applications with an interval of 15 days	250-300 ml/ha
WATERMELON (GREENHOUSE)	4-5 leaves before flowering, fruit setting, fruit ripening periods	100-250 ml/100 L of water	ONIONS	From the period 4-5 leaves to the end of tillering, 2 applications after tillering	300-500 ml/ha
MELON (FIELD)	4-5 leaves before flowering, fruit laying, fruit ripening periods	100-250 ml/100 L of water	OLIVE TREE	The formation of buds before flowering and during the color change of fruits, as well as after harvesting, if necessary.	250-300 ml/100 L of water
APRICOT	Before flowering, when the petals fall off, during the fruiting period, when the fruits reach 6-7 cm in diameter	200-250 ml/100 L of water			



ORGANIC PRODUCTS

LIQUID ORGANIC FERTILIZER OF PLANT ORIGIN

BIODEM

GUARANTEED CONTENT	%W/W
Organic Matter	45
Organic Carbon	20
Total Nitrogen (N)	2
Water-Soluble Potassium Oxide (K ₂ O)	4
pH	4,7-6,7

USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	FOLIAR APPLICATION ml/100 L of water/ha	APPLICATION TO THE SOIL L/ha
TOMATO, EGGPLANT, PEPPER, etc.	- In seedlings (2 applications), - During the period of growth	50-80 80-100	0,8-1 L 0,8-1 L
MELON, WATERMELON, CUCUMBER, etc. (PUMPKIN)	- In the period of early development - Before the first water appears - At the beginning of the sleeve development - During the fruiting period	100-120 100-120 100-120 100-120	0,8-1 L 0,8-1 L 0,8-1 L 0,8-1 L
CHERRY, PEACH, APPLE TREE, PEAR, APRICOT, PLUM, NECTARINE, POMEGRANATE, etc. on deciduous trees	- Immediately after sowing - When the eyes open - After flowering or when the first fruits appear - During the fruit ripening period - After picking fruits	150-250 150-250 150-250 150-250 150-250	10-15 ml per seedling 3-5 years old 15-25 ml 5-10 years old 25-40 ml per tree 10 years and older 40-50 ml per tree 40-50 ml per tree
THE VINEYARD	- Immediately after sowing - When the eyes open - Before flowering - During the corrosion period - After harvesting	150-250 150-250 100-200 100-200 100-200	5 cubes per new root 5-8 cubes per ounce of harvest 0,8-1 L from the dropper 0,8-1 L from drip irrigation 0,8-1 L from drip irrigation
OLIVES	- Immediately after sowing - Before flowering - During the summer period	200-250 200-250 200-250	In June 1-3 or 15-20 ml per tree 3-6 trees 15-25 ml 25-40 ml for 6-12 trees 40-50 ml for 12 or more trees
COTTON	- When 8-10 leaves appear (only on rows of plants) - The period before blooming and flowering - The period of the first cocoon	150-200 150-200 100-150	
CITRUS FRUITS	- Before flowering - After flowering - June Before watering - Immediately after harvest	150-200 150-200 150-200 200-250	Per tree 1-5 years 10-15 ml 4-8 years 15-25 ml 8-15 years old 25-40 ml 40-50 ml over 15 years old
POTATOES, when water is first introduced into the soil, it is given at the rate of 1 liter per 5 decalliters	- The period of the first issue - Before the first water - At the beginning of flowering - After flowering	150-200 150-200 150-200 150-200	The dose for seeds is 100 liters of water per 2 tons of seeds, 100-150 cubes are placed and applied. Spraying is applied 100-200 ml to the decol.
CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND LETTUCE	- After 15-20 days in the period of early development (2-3 ties are made)	150-200 150-200	When planting seedlings, they are dipped in 100 liters of water with a volume of 100-150 ml and planted. There is 0,8-1 L of soil per decalliter.
STRAWBERRY	- At the end of the winter period - Before and after flowering - With an interval of 15 days during harvest	100-150 100-150 100-150	0,8-1 L/ha 2 applications
CUT FLOWERS	- Rose, at the beginning of the waking period - Bud germination - With an interval of 20 days during the development period	150-200 100-150 100-150	0,8-1 L/ha 3-4 applications
GREENHOUSES	- 2 applications during development periods	150-200	Apply 0,8-1 L 2-3 times per 1000 m ² in a greenhouse by sprinkling. Pour clean water from above.

RECOMMENDATIONS

- Can be applied dropwise or on a leaf during the growing season according to the needs of the plants
- It can be mixed with other plant fertilizing preparations and pesticides, with the exception of Bordeaux suspension and alkaline pesticides



**ORGANIC PRODUCTS
LIQUID HUMIC ACID**

FLUX

GUARANTEED CONTENT	%W/W
Organic Matter	%45
Organic Carbon	%20
Total Nitrogen (N)	%2
Water-Soluble Potassium Oxide (K ₂ O)	%4
pH	4-6

ATTENTION

- Keep away from children and food. When applying fertilizers, wear a mask, gloves and safety glasses.
- Never exceed the appropriate dosage and use only if necessary.
- Avoid contact with eyes, face and skin, in case of contact, rinse with plenty of water.
- Do not use used empty packages for other purposes and dispose of them.
- It is recommended to use the fertilizer after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	APPLICATION AND DOSAGE		APPLICATION PERIOD
VEGETABLES (TOMATO, PEPPER, CUCUMBER, EGGPLANT)	SOIL IRRIGATION	3-4 L/ha	After the formation of 3-4 leaves before harvest.
	LEAF WATERING	200-300 ml/100 L of water	
	SOIL IRRIGATION	3000-4000 ml/ha	It begins before flowering and 10 days before harvest.
FRUIT TREES (OLIVE, PISTACHIO, ALMOND, GRAPE, POMEGRANATE, APPLE TREE, PEAR, CHERRY, CHERRY, PEACH, etc.)	LEAF WATERING	200-300 ml/100 L of water	
	SOIL IRRIGATION	2-3 L/ha	After the formation of 3-5 leaves, with an interval of 15 days before harvest.
FIELD CROPS (COTTON, CORN, SOYBEANS, PEANUTS, PADDY, SUNFLOWER, TOBACCO, WHEAT)	LEAF WATERING	200-300 ml/100 L of water	
	SOIL IRRIGATION	1-2 L/ha	With an interval of 15 days during the growing season.
GREEN SPACES, CUT FLOWER CULTURE	LEAF WATERING	200-300 ml/100 L of water	
	SOIL IRRIGATION	2-3 L/ha	With an interval of 15 days during the growing season.
CITRUS FRUITS, BANANAS	LEAF WATERING	200-300 ml/100 L of water	
	SOIL IRRIGATION	3-4 L/ha	With the first top dressing with an interval of 10 days.
MELON, WATERMELON, STRAWBERRY	LEAF WATERING	300-400 ml/100 L of water	



EU FERTILIZERS
BORON(B), MANGO (Mn-Sulfate) and ZINC (Zn-Sulfate)
LIQUID NUTRIENT MIXTURE FOR MICRO PLANTS

AXIS-MICRO

GUARANTEED CONTENT	%W/W
Water-Soluble Boron (B)	5
Water-Soluble Manganese (Mn)	3
Water-Soluble Zinc (Zn)	5

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	LEAF WATERING)	SOIL IRRIGATION
VEGETABLES (open ground) TOMATO, PEPPER, CUCUMBER, EGGPLANT, etc.	It should be used during the formation of color pigments in the product. It significantly improves coloring, attractiveness and quality.	200-300 ml /100 L of water	1-2 L/ha. Apply 1-2 times
APPLE, PEAR, PLUM, APRICOT, CHERRY, PEACH, etc.	It should be used during the formation of color pigments in fruits. Coloring significantly affects attractiveness and quality.	200-300 ml /100 L of water	2-3 L/ha. It should be used 1-2 times
STRAWBERRY	At the coloring stage	200-300 ml /100 L of water	1-2 L/ha
OLIVE	At the coloring stage	200-300 ml /100 L of water	2-3 L/ha
THE VINEYARD	After flowering	200-300 ml /100 L of water	2-3 L/ha
Corn	When forming cobs	200-300 ml /100 L of water	1-2 L/ha
MELON, WATERMELON	It should be used 15-20 days before harvest.	200-300 ml /100 L of water	2-3 L/ha
SUGAR BEET	It should be used 15-20 days before harvest.	200-300 ml /100 L of water	1-2 L/ha
CLOVER	During the growth period	200-300 ml /100 L of water	1-2 L/ha
CUT FLOWERS	During the growth period	200-300 ml /100 L of water	1-2 L/ha
WHEAT, RICE, COTTON, SUNFLOWER	If necessary during the growth period	200-300 ml /100 L of water	1-2 L/ha
TOBACCO	During the growth period	200-300 ml /100 L of water	1-2 L/ha
BANANA	Apply 1-2 times 30 days before harvest.	200-300 ml /100 L of water	1-2 L/ha

RECOMMENDATIONS

- When applying fertilizers, wear a mask, gloves and safety glasses.
- Never exceed the appropriate dose.
- Use only where necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse them with plenty of water.
- After use, wash your hands with plenty of water and dispose of used empty packaging for other purposes.
- It is recommended to use fertilizer after analyzing the soil and/or leaves.



PHOSPO-N

GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	5
Nitrate Nitrogen (NO ₃ -N)	2
Ammonium Nitrogen (NH ₄ -N)	3
Water-Soluble Phosphorus Oxide (P ₂ O ₅)	30
Water-Soluble Zinc (Zn)	4

ATTENTION

- Be careful! Keep away from children and food.
- Read the label before use.
- Do not use used empty packaging for other purposes and dispose of it.
- Wear protective gloves and a mask.
- Wash your hands with plenty of water after use.
- Do not swallow. If swallowed, consult a doctor.
- Avoid contact with skin and eyes, in case of contact, rinse with plenty of water.
- It is recommended to use the fertilizer after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	Type of Applications	
		LEAF WATERING	SOIL IRRIGATION
GREENHOUSE VEGETABLES	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
OUTDOOR VEGETABLES	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
MELON, WATERMELON, STRAWBERRY	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
APPLE, PEAR, QUINCE	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
PEACH, CHERRY, CHERRY, APRICOT, NECTARINE, PLUM	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
GRAPES, BANANA, POMEGRANATE, FIG	Apply 3-4 times with an interval of 20 days After flowering	200-250 ml/100 L of water	1-2 L/ha
CITRUS FRUITS, OLIVE, TEA	Apply 3-4 times with an interval of 20 days After flowering	200-250 ml/100 L of water	1-2 L/ha
HAZELNUT, WALNUT, PISTACHIOS, CHESTNUTS	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
COTTON, CORN, SUNFLOWER, SOY, CANOLA	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
CABBAGE, RADISH, CARROT, CELERY, CAULIFLOWER	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
CEREALS, LEGUMES, FORAGE CROPS	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
SUGAR BEET, POTATOES, ONIONS, GARLIC	Apply 1-2 times from the moment of tubers formation to harvest	200-250 ml/100 L of water	1-2 L/ha
CUT FLOWERS, GREENERY, FIELDS, PADDY	It is applied 2-3 times with an interval of 30 days during the development period	200-250 ml/100 L of water	1-2 L/ha



NEW FOS

GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	5
Nitrate Nitrogen (NO ₃ -N)	2,5
Ammonium Nitrogen (NH ₄ -N)	2,5
Water-Soluble Phosphorus Pentoxide (P ₂ O ₅)	20

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	METHOD OF APPLICATION	
		LEAF WATERING	SOIL IRRIGATION
FOR GREENHOUSE VEGETABLES	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
FOR VEGETABLES IN THE OPEN GROUND	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
MELON, WATERMELON, STRAWBERRY	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
APPLE, PEAR, QUINCE	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
PEACH, CHERRY, CHERRY, APRICOT, NECTARINE, PLUM	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
GRAPES, BANANA, POMEGRANATE, FIG	Apply 3-4 times with an interval of 20 days After flowering	200-250 ml/100 L of water	1-2 L/ha
CITRUS FRUITS, OLIVES, TEA	Apply 3-4 times with an interval of 20 days After flowering	200-250 ml/100 L of water	1-2 L/ha
HAZELNUT, WALNUTS, PISTACHIOS, PISTACHIOS, CHESTNUTS	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
COTTON, CORN, SUNFLOWER, SOY, CANOLA	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
CABBAGE, RADISH, CARROT, CELERY, CAULIFLOWER	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
CEREALS, LEGUMES, FEED, AGRICULTURAL CROPS	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
SUGAR BEET, POTATOES, ONIONS, GARLIC	Apply 1-2 times from the moment of tubers formation to harvest	200-250 ml/100 L of water	1-2 L/ha
FOR CUT FLOWERS, GREEN-ERY, FIELDS, RICE	It is applied 2-3 times with an interval of 30 days during the development period	200-250 ml/100 L of water	1-2 L/ha

ATTENTION

- Keep away from children and food.
- Read the label before use.
- Do not use used empty packages for other purposes and do not destroy them.
- Use protective gloves and a mask.
- Wash your hands with plenty of water after use.
- Do not swallow. If swallowed, consult a doctor.
- Avoid contact with skin and eyes, in case of contact, rinse with plenty of water.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.



SOLUTION CALCIUM CHLORIDE FOR FERTILIZERS

ONE PLUS

GUARANTEED CONTENT	%W/W
Water-Soluble Calcium Oxide (CaO)	14
Water-Soluble Boron (B)	0,3

ATTENTION

- Never exceed the appropriate dose. Use it only if necessary.
- Keep away from children and food.
- In case of contact with eyes, face and skin, rinse with plenty of water.
- After use, wash your hands with plenty of water and dispose of used empty packages for other purposes.
- When applying fertilizers, use a mask, gloves and glasses.
- It is recommended to use the fertilizer after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	DOSES AND TIME OF APPLICATION	
	LEAF WATERING	SOIL IRRIGATION
VEGETABLES IN THE OPEN GROUND	250-300 ml / 100 L of water	400-500 ml/ha
IN GREENHOUSES IN THE COOL HOURS OF THE DAY	150-250 ml / 100 L of water	350-400 ml/ha
SOFT STONE FRUITS	300-350 ml / 100 L of water	450-600 ml or 50 ml on the tree
HARD STONE FRUITS	300-350 ml / 100 L of water	450-500 ml or 50 ml on the tree
THE VINEYARD	250-300 ml / 100 L of water	300-400 ml or 50 ml on the tree
CITRUS FRUITS, OLIVE TREES	250-300 ml / 100 L of water	400-450 ml or 50 ml on the tree



SOLUTION OF ZINC FERTILIZERS (CONTAINS ZINC SULFATE)

ZINCOMİN

GUARANTEED CONTENT	%W/W
Water-Soluble Zinc (Zn)	11

DOES NOT CONTAIN CHLORINE

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	METHOD OF APPLICATION	
		LEAF WATERING	SOIL IRRIGATION
FOR GREENHOUSE VEGETABLES	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
FOR VEGETABLES IN THE OPEN GROUND	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
MELON, WATERMELON, STRAWBERRY	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
APPLE, PEAR, QUINCE	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
PEACH, CHERRY, CHERRY, APRICOT, NECTARINE, PLUM	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
GRAPES, BANANA, POMEGRANATE, FIG	Apply 3-4 times with an interval of 20 days After flowering	200-250 ml/100 L of water	1-2 L/ha
CITRUS FRUITS, OLIVES, TEA	Apply 3-4 times with an interval of 20 days After flowering	200-250 ml/100 L of water	1-2 L/ha
HAZELNUT, WALNUTS, PISTACHIOS, PISTACHIOS, CHESTNUTS	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
COTTON, CORN, SUNFLOWER, SOY, CANOLA	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
CABBAGE, RADISH, CARROT, CELERY, CAULIFLOWER	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
CEREALS, LEGUMES, FEED, AGRICULTURAL CROPS	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
SUGAR BEET, POTATOES, ONIONS, GARLIC	Apply 1-2 times from the moment of tubers formation to harvest	200-250 ml/100 L of water	1-2 L/ha
FOR CUT FLOWERS, GREENERY, FIELDS, RICE	It is applied 2-3 times with an interval of 30 days during the development period	200-250 ml/100 L of water	1-2 L/ha

ATTENTION

- Keep away from children and food.
- Do not exceed the dosage and use only if necessary.
- Avoid contact with eyes, face and skin, in case of contact, rinse with plenty of water.
- Do not use used empty packages for other purposes and destroy them.
- It is recommended to use the fertilizer after analyzing the soil and/or leaves.



REDOX CU

GUARANTEED CONTENT	%W/W
Water-Soluble Copper (Cu)	5

ATTENTION

- Wear a mask, gloves and safety glasses during fertilization.
- Never exceed the appropriate dose.
- Use only where necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse them with plenty of water.
- After use, wash your hands with plenty of water and dispose of used empty packaging for other purposes.
- It is recommended to use the fertilizer after analyzing the soil and/or leaves.



USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	METHOD OF APPLICATION	
		LEAF WATERING	SOIL IRRIGATION
GREENHOUSE VEGETABLES	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
OUTDOOR VEGETABLES	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
MELON, WATERMELON, STRAWBERRY	Apply 4-5 times from the moment 4-5 leaves appear on plants until harvest	200-250 ml/100 L of water	1-2 L/ha
APPLE, PEAR, QUINCE	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
PEACH, CHERRY, CHERRY, APRICOT, NECTARINE, PLUM	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
GRAPES, BANANA, POMEGRANATE, FIG	Apply 3-4 times with an interval of 20 days After flowering	200-250 ml/100 L of water	1-2 L/ha
CITRUS FRUITS, OLIVE, TEA	Apply 3-4 times with an interval of 20 days After flowering	200-250 ml/100 L of water	1-2 L/ha
HAZELNUT, WALNUT, PISTACHIOS, CHESTNUTS	It is applied 3-4 times with an interval of 20 days from the beginning of fruiting.	200-250 ml/100 L of water	1-2 L/ha
COTTON, CORN, SUNFLOWER, SOY, CANOLA	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
CABBAGE, RADISH, CARROT, CELERY, CAULIFLOWER	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
CEREALS, LEGUMES, FORAGE CROPS	Apply 1-2 times from the moment 4-5 leaves appear on plants to harvest	200-250 ml/100 L of water	1-2 L/ha
SUGAR BEET, POTATOES, ONIONS, GARLIC	Apply 1-2 times from the moment of tubers formation to harvest	200-250 ml/100 L of water	1-2 L/ha
CUT FLOWERS, GREEN SPACES, PADDY	During the development period, it is applied 2-3 times with an interval of 30 days	200-250 ml/100 L of water	1-2 L/ha

ORGANIC PRODUCTS LIQUID ORGANIC FERTILIZER OF PLANT ORIGIN

REDOX B

GUARANTEED CONTENT

%W/W

Water-Soluble Boron (B)

8

ATTENTION

- Never exceed the appropriate dose.
- Use only if necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse with plenty of water.
- After use, wash your hands with plenty of water and dispose of used empty packaging for other purposes.
- Wear a mask, gloves and safety glasses during fertilization.

USE RECOMMENDATIONS

CROP	LEAF WATERING	SOIL IRRIGATION
OUTDOOR VEGETABLES	150-180 ml / 100 L of water	200-300 ml/ha
GREENHOUSES	100-150 ml / 100 L of water	250-300 ml/ha
SOFT STONE FRUITS	100-150 ml / 100 L of water	200-300 ml/ha or 50 ml per tree
HARD STONE FRUITS	100-150 ml / 100 L of water	250-300 ml/ha or 50 ml per tree
VINEYARD	150-200 ml / 100 L of water	200-250 ml/ha or 50 ml per tree
CITRUS FRUITS, OLIVE TREES	100-150 ml / 100 L of water	200-250 ml/ha or 50 ml per tree
WATERMELON, CAULIFLOWER, ONION, GARLIC, etc.	100-150 ml / 100 L of water	250-300 ml/ha
COTTON, TOBACCO, BEET, CORN etc.	100-150 ml / 100 L of water	250-300 ml/ha
WHEAT, BARLEY, PADDY etc.	100-150 ml / 100 L of water	250-300 ml/ha



NEWAMIN

GUARANTEED CONTENT	%W/W
Total Amount Of Organic Matter	35
Organic Carbon	12
Total Nitrogen (N)	2
Water-Soluble Potassium Oxide (K ₂ O)	2
Free Amino Acids	11
pH	6,7-8,7

ATTENTION

- Be careful! Keep away from children and food.
- Read the label before use.
- Do not use used empty packaging for other purposes and dispose of it.
- Use protective gloves and a mask.
- Wash your hands with plenty of water after use.
- Do not swallow. If swallowed, consult a doctor.
- Avoid contact with skin and eyes, in case of contact, rinse with plenty of water.
- It is recommended to use the fertilizer after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	FOLIAR APPLICATION ml/100 L of water/ha	APPLICATION TO THE SOIL L/ha
TOMATO, EGGPLANT, PEPPER, etc.	- In seedlings (2 applications), - During the period of growth	50-80 80-100	0,8-1 L 0,8-1 L
MELON, WATERMELON, CUCUMBER, etc. (PUMPKIN)	- In the period of early development - Before the first water appears - At the beginning of the sleeve development - During the fruiting period	100-120 100-120 100-120 100-120	0,8-1 L 0,8-1 L 0,8-1 L 0,8-1 L
CHERRY, PEACH, APPLE TREE, PEAR, APRICOT, PLUM, NECTARINE, POME- granate, etc. on deciduous trees	- Immediately after sowing - When the eyes open - After flowering or when the first fruits appear - During the fruit ripening period - After picking fruits	150-250 150-250 150-250 150-250 150-250	10-15 ml per seedling 3-5 years old 15-25 ml 5-10 years old 25-40 ml per tree 10 years and older 40-50 ml per tree 40-50 ml per tree
THE VINEYARD	- Immediately after sowing - When the eyes open - Before flowering - During the corrosion period - After harvesting	150-250 150-250 100-200 100-200 100-200	5 cubes per new root 5-8 cubes per ounce of harvest 0,8-1 L from the dropper 0,8-1 L from drip irrigation 0,8-1 L from drip irrigation
OLIVES	- Immediately after sowing - Before flowering - During the summer period	200-250 200-250 200-250	In June 1-3 or 15-20 ml per tree 3-6 trees 15-25 ml 25-40 ml for 6-12 trees 40-50 ml for 12 or more trees
COTTON	- When 8-10 leaves appear (only on rows of plants) - The period before blooming and flowering - The period of the first cocoon	150-200 150-200 100-150	
CITRUS FRUITS	- Before flowering - After flowering - June Before watering - Immediately after harvest	150-200 150-200 150-200 200-250	Per tree 1-5 years 10-15 ml 4-8 years 15-25 ml 8-15 years old 25-40 ml 40-50 ml over 15 years old
POTATOES, when water is first introduced into the soil, it is given at the rate of 1 liter per 5 decaliters	- The period of the first issue - Before the first water - At the beginning of flowering - After flowering	150-200 150-200 150-200 150-200	The dose for seeds is 100 liters of water per 2 tons of seeds, 100, 150 cubes are placed and applied. Spraying is applied 100-200 ml to the decar.
CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND LETTUCE	- After 15-20 days in the period of early development (2-3 ties are made)	150-200 150-200	When planting seedlings, they are dipped in 100 liters of water with a volume of 100-150 ml and planted. There is 0,8-1 L of soil per decaliter.
STRAWBERRY	- At the end of the winter period - Before and after flowering - With an interval of 15 days during harvest	100-150 100-150 100-150	0,8-1 L/ha 2 applications
CUT FLOWERS	- Rose, at the beginning of the waking period - Bud germination - With an interval of 20 days during the development period	150-200 100-150 100-150	0,8-1 L/ha 3-4 applications
GREENHOUSES	- 2 applications during development periods	150-200	Apply 0,8-1 L 2-3 times per 1000 m ² in a greenhouse by sprinkling. Pour clean water from above.



PRODUCTS FROM ORGANIC PLANTS
LIQUID SEAWEED

MERMAID

GUARANTEED CONTENT	%W/W
Organic Matter	15
Alginic Acid	0,3
Water-Soluble Potassium Oxide (K ₂ O)	3
Maximum EC (dS/m)	15
pH	4,7-6,7

USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	LEAF WATERING	DRIP IRRIGATION
VEGETABLES: TOMATOES, PEPPERS, EGGPLANTS, BEANS, PEAS, POTATOES, ONIONS, GARLIC, CUCUMBER, MELON, WATERMELON, PUMPKIN ETC. (ALL TYPES OF OPEN AND CLOSED GROUND)	- In the early stages of plant development (flower tying) - From flowering to harvest (2-3 applications)	100-150 ml 100-150 ml	400-500 ml
FRUIT TREES: SOFT AND STONE FRUITS, CHERRIES, CHERRIES, PEACHES, APRICOTS, PLUMS, CHERRIES, AP- PLE, PEAR, ALMOND, HAZELNUT, POMEGRANATE ETC.	- The beginning of flowering, fruiting 1 application during the period of setting, fruit development and coloring	100-150 ml	400-500 ml
THE VINEYARD: TABLE GRAPES AND RAISINS SEEDLINGS FOR ALL VARIETIES.	- With the growth of shoots 3-5 cm, during the period of cluster release, before flowering, on thin and large bushes, during freshwater walking.	100-150 ml	400-500 ml
STRAWBERRY	During the season	100-150 ml	400-500 ml
COTTON	- from the period with 6-8 leaves to harvest	100-150 ml	400-500 ml
GREENHOUSE AND CUT FLOWERS	- Throughout the entire period of development	100-150 ml	400-500 ml
LETTUCE, PARSLEY, WATERCRESS, ARUGULA etc. (GREEN PLANTS)	- Throughout the entire period of development	100-150 ml	400-500 ml
FIELD CROPS: WHEAT, BARLEY, OATS, RICE etc.	2 applications with an interval of 10 days during the tillering period.	100-150 ml	400-500 ml
OLIVES	- The beginning of flowering, fruiting Up to the period of tying and coloring, throug- hout the summer period.	100-150 ml	400-500 ml
CITRUS FRUITS	- The flowering period, the period of fruit setting and before harvest.	100-150 ml	400-500 ml
TOBACCO	- 3 applications during the seedling period, from plant- ing to thinning period.	100-150 ml	400-500 ml
INDUSTRIAL CROPS: SUGAR BEET, ANISE, CORN, SUNFLOWER etc.	- At least 2 applications during the early development period	100-150 ml	400-500 ml
CHICKPEAS	- Minimum of 2 applications at the beginning of flowering	100-150 ml	400-500 ml

RECOMMENDATIONS

- Can be applied drip or on a leaf during the growing season, depending on the needs of the plants.
- It can be mixed with other plant nutrition products and pesticides, with the exception of Bordeaux slurry and alkaline pesticides.



GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	3
Urea Nitrogen (NH ₂ -N)	3
Water-Soluble Potassium Oxide (K ₂ O)	25

ATTENTION

- Never exceed the appropriate dose.
- Use only if necessary.
- Keep away from children and food.
- CAUSES IRRITATION. Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse them with plenty of water.
- After use, wash your hands with plenty of water and dispose of used empty packaging for other purposes.
- Wear a mask, gloves and safety glasses during fertilization.

USE RECOMMENDATIONS

CROP	LEAF WATERING	DRIP IRRIGATION
OUTDOOR VEGETABLES	200-300 ml / 100 L of water	1-2 L/ha
GREENHOUSES	150-250 ml / 100 L of water	1,5-2 L/ha
SOFT STONE FRUITS	300-350 ml / 100 L of water	2-3 L/ha or 100 g per tree
HARD STONE FRUITS	300-350 ml / 100 L of water	1-2 L/ha or 70 g per tree
VINEYARD	250-300 ml / 100 L of water	1,5-2 L/ha or 60 g per tree
CITRUS FRUITS, OLIVE TREES	250-300 ml / 100 L of water	1-2 L/ha or 70 g per tree
WATERMELON, CAULIFLOWER, ONION, GARLIC, etc.	150-200 ml / 100 L of water	1,5-2 L/ha
COTTON, TOBACCO, BEET, CORN etc.	250-300 ml / 100 L of water	1,5-2 L/ha
WHEAT, BARLEY, PADDY etc.	150-250 ml / 100 L of water	1-2 L/ha



EC FERTILIZER NPK FERTILIZER SOLUTION

TETRA

RECOMMENDATIONS

- Keep away from children and food. Use a mask, gloves and glasses during fertilization.
- Avoid contact with eyes, face and skin. In case of contact, rinse with plenty of water.
- Dispose of used empty packaging and do not use it for other purposes.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.

TETRA 7-7-7

GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	7
Urea Nitrogen (N)	7
Water-Soluble Phosphorus Pentoxide (P ₂ O ₅)	7
Water-Soluble Potassium Oxide (K ₂ O)	7

TETRA 4-6-10

GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	4
Urea Nitrogen (N)	4
Water-Soluble Phosphorus Pentoxide (P ₂ O ₅)	6
Water-Soluble Potassium Oxide (K ₂ O)	10

TETRA 7-21-3

GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	7
Ammonium Nitrogen (N)	4
Nitrate Nitrogen (N)	3
Water-Soluble Phosphorus Pentoxide (P ₂ O ₅)	21
Water-Soluble Potassium Oxide (K ₂ O)	3

USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	LEAF WATERING	DRIP IRRIGATION
VEGETABLES IN THE OPEN GROUND	After transplanting the seedlings, the treatment is carried out every 20 days throughout the season.	200-250 ml	450-500 ml/day
IN GREENHOUSES	In the form of weekly treatments, starting from the seedling period.	150-250 ml	350-400 ml/day
IN FRUITS WITH SOFT SEEDS	The application is made every 20 days During the season, starting from the seedling stage.	200-250 ml	450-600 ml/day or 50 ml per tree
IN STONE FRUITS	After the pharyngeal period, every 25 days.	250-300 ml	450-600 ml/day or 50 ml per tree
VINEYARD	The treatment is carried out every 25 days After flowering throughout the season.	250-300 ml	300-400 ml/day or 50 ml per tree
CITRUS FRUITS OR OLIVE TREES	The application is made every 20 days During the season, starting from the seedling stage.	250-300 ml	400-450 ml/day or 50 ml per tree
WATERMELON, CAULIFLOWER, ONION, GARLIC	It is applied throughout the season with an interval of 15 days after the second weeding.	150-250 ml	350-400 ml/day
COTTON, TOBACCO, SWEET BEET, CORN etc.	It is applied throughout the season with an interval of 15 days after the second weeding.	150-250 ml	350-400 ml/day
WHEAT, BARLEY, RICE etc.	It is applied throughout the season with an interval of 15 days after the second weeding.	150-250 ml	350-400 ml/day



EU FERTILIZERS
A MIXTURE OF MICRONUTRIENT PLANTS OF MANGANESE (MN-SULFATE)
AND ZINC (ZN-SULFATE)

EXE MNZN

GUARANTEED CONTENT	%W/W
Water-Soluble Manganese (Mn)	14
Water-Soluble Zinc (Zn)	12

RECOMMENDATIONS

- Use a mask, gloves and glasses during fertilization.
- Never exceed the appropriate dose.
- Use only when necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse with plenty of water.
- After use, wash your hands with plenty of water and discard the used empty packaging before using it for other purposes.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	LEAF WATERING	DRIP IRRIGATION
VEGETABLES (OPEN GROUND), TOMATOES, PEPPERS, CUCUMBERS, Eggplants etc.	It should be used in the formation of color pigments in the product. Coloring significantly affects the appearance and quality.	200-300g/100 L of water	1-2 kg/ha. It should be applied 1-2 times.
APPLE, PEAR, PLUM, APRICOT, CHERRY, PEACH etc.	It should be used in the formation of color pigments in fruits. Coloring significantly affects attractiveness and quality.	200-300g/100 L of water	2-3 kg/ha. It should be applied 1-2 times.
STRAWBERRY	At the stage of staining.	200-300g/100 L of water	1-2 kg/ha
OLIVE	At the stage of staining.	200-300g/100 L of water	2-3 kg/ha
THE VINEYARD	At the stage of green grapes.	200-300g/100 L of water	2-3 kg/ha
CORN	In the cob formation.	200-300g/100 L of water	1-2 kg/ha
MELON, WATERMELON	It should be used 15-20 days before harvest.	200-300g/100 L of water	2-3 kg/ha
SUGAR BEET	It should be used 15-20 days before harvest.	200-300g/100 L of water	1-2 kg/ha
CLOVER	During the period of growth	200-300g/100 L of water	1-2 kg/ha
CUT FLOWERS	During the period of growth	200-300g/100 L of water	1-2 kg/ha
WHEAT, RICE, COTTON, SUNFLOWER	If necessary during the growth period	200-300g/100 L of water	1-2 kg/ha
TOBACCO	During the period of growth	200-300g/100 L of water	1-2 kg/ha
BANANA	Apply 1-2 times 30 days before harvest.	200-300g/100 L of water	1-2 kg/ha



EU FERTILIZERS
NPK FERTILIZER SOLUTION

SET-UP

RECOMMENDATIONS

Water-soluble metal salts turn into metal salts, such as metal oxides-metal carbonates, which are insoluble in water and therefore cannot be used by plants if the pH of the soil and water are high.

SET UP has a good ability to retain cations due to its complexing ability. Under all conditions, it forms a complex structure with compounds of metals and alkaline earth metals, such as oxides, hydroxides and carbonates, and binds to itself, forming a complex of water-soluble metal salts.

creates.

This complex structure, which the plant easily assimilates, is stable. Carbonate and bicarbonate salts of calcium and magnesium, which are present in the structure of irrigation water and cause temporary and permanent stiffness, form a structure in pipes that we call calcification. For this reason, blockages occur in irrigation pipes from time to time.

SET UP is used to prevent these locks. SET UP is not completely acidic and has a suitable pH level. Thanks to this function, it is an ideal pH regulator for irrigation water.

It does not contain mineral acids such as nitric and phosphoric acids, which can cause soil salinization.



EU FERTILIZERS
A MIXTURE OF SOLID MICRO-PLANT NUTRIENTS BORON (B),
COPPER (CU-SULFATE), IRON (FE-SULFATE), MANGANESE (MN-SUL-
FATE) AND ZINC (ZN-SULFATE)

MIKROFIX COMBI

GUARANTEED CONTENT	%W/W
Water-Soluble Boron (B)	1
Water-Soluble Copper (Cu)	0,5
Water-Soluble Iron (Fe)	4
Water-Soluble Manganese (Mn)	4
Water-Soluble Zinc (Zn)	4

RECOMMENDATIONS

- Use a mask, gloves and glasses during fertilization.
- Never exceed the appropriate dose.
- Use only when necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse with plenty of water.
- After use, wash your hands with plenty of water and discard the used empty packaging before using it for other purposes.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	APPLICATION
VEGETABLES, TOMATOES, PEPPERS, EGGPLANTS ETC.	It is applied 2-3 weeks apart at the beginning of flowering.	100-150 g/100 L of water
FRUITS, CITRUS FRUITS, APPLES, PEARS ETC.	It is used when the fruits reach the size of a walnut and on the lower shoots.	100-150 g/100 L of water
VINEYARD	2-3 applications, starting from the ripening of the bunch	10-20 g / grapevine
CORN	It is used when the plant reaches 40-50 cm in height.	150 g/ha
COTTON	It is used immediately before and immediately After flowering.	100-200 g/ha
SUGAR BEET	It is used in the 6-8-leaf period.	100 g/ha
POTATO	During the formation of tubers, 2-3 treatments are performed.	150 g/ha
ONION	It is applied twice 30 days after planting.	100 g/ha
WHEAT	The application is made during the tillering period.	100 g/ha
CUT FLOWERS	From the very beginning, 2-3 applications are made.	100 g/100 L of water
MELON, WATERMELON	It is applied 2-3 times from the moment of fruit formation.	100 g/ha



ORGANIC PRODUCTS LIQUID HUMIC ACID

HUMATİN

GUARANTEED CONTENT	%W/W
Organic Matter	12
Total (Humic + Fulvic Acid)	15
pH	9,3-11,3

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	METHOD OF APPLICATION	
		LEAF WATERING	SOIL IRRIGATION
CITRUS FRUITS	* Before flowering * When the fruits reach large sizes	200-250 ml/100 L of water	1-3 L/ha
APPLE, PEAR etc.	* Before flowering and After flowering.	200-250 ml/100 L of water	1-3 L/ha
PEACH, APRICOT, CHERRY, CHERRY, PLUM AND POMEGRANATE	* Before flowering and After flowering.	200-250 ml/100 L of water	1-3 L/ha
VINEYARD	* Before flowering * When the fruits reach large sizes	200-250 ml/100 L of water	1-3 L/ha
PISTACHIO, HAZELNUT	* Before flowering * After the fruit is tied	200-250 ml/100 L of water	1-3 L/ha
OLIVE	* Before flowering * When the fruits reach large sizes	200-250 ml/100 L of water	1-3 L/ha
BANANA	* Before flowering * During the period of fruit growth	200-250 ml/100 L of water	1-3 L/ha
KIWI, FIGS	* Before flowering * During the period of fruit growth	200-250 ml/100 L of water	1-3 L/ha
TOMATOES, PEPPERS	* Before flowering * During the period of fruit growth	200-250 ml/100 L of water	1-3 L/ha
CUCUMBER, EGGPLANT	* Before flowering * During the period of fruit growth	200-250 ml/100 L of water	1-3 L/ha
MELON, WATERMELON	* 10 days after landing * Before flowering	200-250 ml/100 L of water	1-3 L/ha
CUT FLOWERS	* Before flowering * During the period of fruit growth	200-250 ml/100 L of water	1-3 L/ha
LENTILS, CHICKPEAS	* Before flowering * During the period of fruit growth	50 ml – 1 L/day	1-3 L/ha
PEANUT	* Before flowering * During the period of fruit growth	500 ml – 1 L/day	1-3 L/ha
COTTON, SUNFLOWERS, CORN, CANOLA, etc.	* When it reaches a height of 20-40 cm. * When it reaches a height of 50-70 cm.	200-250 ml/day	1-3 L/ha

RECOMMENDATIONS

- It can be applied without drops or on leaves during the growing season, depending on the needs of the plants
- Can be mixed with other plant foods and pesticides, with the exception of Bordeaux suspension and alkaline preparations



HEADLAND COOPER

GUARANTEED CONTENT	%W/W
Total Copper (Cu)	19

RECOMMENDATIONS

- Use a mask, gloves and glasses during fertilization.
- Never exceed the appropriate dose.
- Use only when necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse with plenty of water.
- After use, wash your hands with plenty of water and discard the used empty packaging before using it for other purposes.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	SOIL IRRIGATION
FOR GREENHOUSE VEGETABLES	It is applied 4-9 times from stage 4-5 leaves of plants to harvest.	1-2 L/ha
FOR VEGETABLES IN THE OPEN GROUND	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	1-2 L/ha
MELON, WATERMELON, STRAWBERRY	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	1-2 L/ha
APPLE, PEAR, QUINCE	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	1-2 L/ha
PEACH, CHERRY, APRICOT, NECTARINE, PLUM	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	1-2 L/ha
GRAPES, BANANA, POMEGRANATE, FIG	Apply 3-4 times with an interval of 20 days After flowering.	1-2 L/ha
CITRUS FRUITS, OLIVES, TEA	Apply 3-4 times with an interval of 20 days After flowering.	1-2 L/ha
HAZELNUT, WALNUTS, PISTACHIOS, CHESTNUTS	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	1-2 L/ha
COTTON, CORN, SUNFLOWER, SOY, CANOLA	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	1-2 L/ha
CABBAGE, RADISH, CARROT, CELERY, KAMABAHAR	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	1-2 L/ha
CEREALS, LEGUMES, FORAGE CROPS	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	1-2 L/ha
SUGAR BEET, POTATOES, ONIONS, GARLIC	Apply 1-2 times from the formation of tubers to harvest.	1-2 L/ha
CUT FLOWERS, GREEN SPACES, RICE	It is applied 2-3 times with an interval of 30 days during the development period.	1-2 L/ha



EU FERTILIZERS
NK FERTILIZER SOLUTION

HEADLAND HI-K

GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	3
Urea Nitrogen (N)	3
Water-Soluble Potassium Oxide (K ₂ O)	28

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	METHOD OF APPLICATION	
		LEAF WATERING	SOIL IRRIGATION
FOR GREENHOUSE VEGETABLES	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
FOR VEGETABLES IN THE OPEN GROUND	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
MELON, WATERMELON, STRAWBERRY	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
APPLE, PEAR, QUINCE	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
PEACH, CHERRY, CHERRY, APRICOT, NECTARINE, PLUM	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
GRAPES, BANANAS, POMEGRANATES, FIGS	Apply 3-4 times with an interval of 20 days After flowering.	200-250 ml / 100 L of water	1-2 L/ha
CITRUS FRUITS, OLIVES, TEA	Apply 3-4 times with an interval of 20 days After flowering.	200-250 ml / 100 L of water	1-2 L/ha
HAZELNUT, WALNUTS, PISTACHIOS, CHESTNUTS	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
COTTON, CORN, SUNFLOWER, SOY, CANOLA	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
CABBAGE, RADISH, CARROT, CELERY, CAULIFLOWER	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
CEREALS, LEGUMES, FORAGE CROPS	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
SUGAR BEET, POTATOES, ONIONS, GARLIC	Apply 1-2 times from the formation of tubers to harvest.	200-250 ml / 100 L of water	1-2 L/ha
CUT FLOWERS, GREEN SPACES, RICE FIELDS	It is applied 2-3 times with an interval of 30 days during the development period.	200-250 ml / 100 L of water	1-2 L/ha

RECOMMENDATIONS

- Use a mask, gloves and glasses during fertilization.
- Never exceed the appropriate dose.
- Use only when necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse with plenty of water.
- After use, wash your hands with plenty of water and discard the used empty packaging before using it for other purposes.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.



EC FERTILIZERS
A SOLUTION OF NITROGEN FERTILIZERS

HEADLAND HI-N

GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	24
Ammonium Nitrogen (NH ₄ -N)	6
Nitrate Nitrogen (N)	6
Urea Nitrogen (NH ₃ -N)	12
Water-Soluble Copper (Cu)	0,6
Water-Soluble Manganese (Mn)	0,6

RECOMMENDATIONS

- Use a mask, gloves and glasses during fertilization.
- Never exceed the appropriate dose.
- Use only when necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse with plenty of water.
- After use, wash your hands with plenty of water and discard the used empty packaging before using it for other purposes.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	METHOD OF APPLICATION	
		LEAF WATERING	SOIL IRRIGATION
FOR GREENHOUSE VEGETABLES	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
FOR VEGETABLES IN THE OPEN GROUND	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
MELON, WATERMELON, STRAWBERRY	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
APPLE, PEAR, QUINCE	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
PEACH, CHERRY, CHERRY, APRICOT, NECTARINE, PLUM	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
GRAPES, BANANAS, POMEGRANATES, FIGS	Apply 3-4 times with an interval of 20 days After flowering.	200-250 ml / 100 L of water	1-2 L/ha
CITRUS FRUITS, OLIVES, TEA	Apply 3-4 times with an interval of 20 days After flowering.	200-250 ml / 100 L of water	1-2 L/ha
HAZELNUT, WALNUTS, PISTACHIOS, CHESTNUTS	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
COTTON, CORN, SUNFLOWER, SOY, CANOLA	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
CABBAGE, RADISH, CARROT, CELERY, CAULIFLOWER	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
CEREALS, LEGUMES, FORAGE CROPS	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
SUGAR BEET, POTATOES, ONIONS, GARLIC	Apply 1-2 times from the formation of tubers to harvest.	200-250 ml / 100 L of water	1-2 L/ha
CUT FLOWERS, GREEN SPACES, RICE FIELDS	It is applied 2-3 times with an interval of 30 days during the development period.	200-250 ml / 100 L of water	1-2 L/ha



EU FERTILIZERS
NK FERTILIZER SOLUTION

HEADLAND HI-PHOS

GUARANTEED CONTENT	%W/W
Water-Soluble Phosphorus Pentoxide (P ₂ O ₅)	10
Water-Soluble Potassium Oxide (K ₂ O)	5,1
Water-Soluble Magnesium Oxide (MgO)	5,5

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	METHOD OF APPLICATION	
		LEAF WATERING	SOIL IRRIGATION
FOR GREENHOUSE VEGETABLES	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
FOR VEGETABLES IN THE OPEN GROUND	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
MELON, WATERMELON, STRAWBERRY	Apply 4-5 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
APPLE, PEAR, QUINCE	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
PEACH, CHERRY, CHERRY, APRICOT, NECTARINE, PLUM	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
GRAPES, BANANAS, POMEGRANATES, FIGS	Apply 3-4 times with an interval of 20 days After flowering.	200-250 ml / 100 L of water	1-2 L/ha
CITRUS FRUITS, OLIVES, TEA	Apply 3-4 times with an interval of 20 days After flowering.	200-250 ml / 100 L of water	1-2 L/ha
HAZELNUT, WALNUTS, PISTACHIOS, CHESTNUTS	Apply 3-4 times with an interval of 20 days, starting with the fruit setting.	200-250 ml / 100 L of water	1-2 L/ha
COTTON, CORN, SUNFLOWER, SOY, CANOLA	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
CABBAGE, RADISH, CARROT, CELERY, CAULIFLOWER	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
CEREALS, LEGUMES, FORAGE CROPS	Apply 1-2 times from the stage 4-5 leaves of plants before harvesting.	200-250 ml / 100 L of water	1-2 L/ha
SUGAR BEET, POTATOES, ONIONS, GARLIC	Apply 1-2 times from the formation of tubers to harvest.	200-250 ml / 100 L of water	1-2 L/ha
CUT FLOWERS, GREEN SPACES, RICE FIELDS	It is applied 2-3 times with an interval of 30 days during the development period.	200-250 ml / 100 L of water	1-2 L/ha

RECOMMENDATIONS

- ATTENTION! IT CAUSES IRRITATION!
- Use a mask, gloves and glasses during fertilization.
- Never exceed the appropriate dose.
- Use only when necessary.
- Keep away from children and food.
- Avoid contact with eyes, face and skin. In case of contact with eyes and hands, rinse with plenty of water. After use, wash your hands with plenty of water and discard the used empty packaging, rather than using it with another package.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.



EU FERTILIZERS
A MIXTURE OF MICRO-PLANT NUTRIENTS
BORA (B) AND ZINC (ZN-SULFATE)

TETRA ZnB

GUARANTEED CONTENT	%W/W
Water-Soluble Boron (B)	10
Water-Soluble Zinc (Zn)	12

RECOMMENDATIONS

- Keep away from children and food.
- Read the label before use.
- Do not use used empty packaging for other purposes and do not destroy it.
- Use protective gloves and a mask.
- Wash your hands with plenty of water after use.
- Do not swallow. If swallowed, consult a doctor.
- Avoid contact with skin and eyes, in case of contact, rinse with plenty of water.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.

USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	LEAF WATERING	DRIP IRRIGATION
VEGETABLES: TOMATOES, PEPPERS, EGGPLANTS, BEANS, PEAS, POTATOES, ONIONS, GARLIC, CUCUMBER, MELON, WATERMELON, PUMPKIN etc. (ALL TYPES OF OPEN AND CLOSED GROUND)	- In the early stages of plant development (flower tying) - From flowering to harvest (2-3 applications)	100-150 g 100-150 g	400-500 g
FRUIT TREES: SOFT AND STONE FRUITS, CHER- RIES, PEACHES, APRICOTS, PLUMS, CHERRIES, APPLE, PEAR, ALMOND, HAZELNUT, POMEGRANATE ETC.	- The beginning of flowering, fruiting 1 application during the period of setting, fruit development and coloring	100-150 g	400-500 g
VITICULTURE: TABLE GRAPES AND RAISINS SEEDLINGS FOR ALL VARIETIES.	- With the growth of shoots 3-5 cm, during the period of cluster release, before flowering, on thin and large bushes, during freshwater walking.	100-150 g	400-500 g
STRAWBERRY	During the season	100-150 g	400-500 g
COTTON	- from the period with 6-8 leaves to harvest	100-150 g	400-500 g
GREENHOUSE AND CUT FLOWERS	- Throughout the entire period of development	100-150 g	400-500 g
LETTUCE, PARSLEY, WATERCRESS, ARUGULA etc. (GREEN PLANTS)	- Throughout the entire period of development	100-150 g	400-500 g
FIELD CROPS: WHEAT, BARLEY, OATS, RICE etc.	2 applications with an interval of 10 days during the tillering period.	100-150 g	400-500 g
OLIVE TREES	- The beginning of flowering, fruiting Up to the period of tying and coloring, throughout the summer period.	100-150 g	400-500 g
CITRUS FRUITS	- The flowering period, the period of fruit setting and before harvest.	100-150 g	400-500 g
TOBACCO	- 3 applications during the seedling period, from plant- ing to thinning period.	100-150 g	400-500 g
INDUSTRIAL CROPS: SUGAR BEET, ANISE, CORN, SUNFLOWER ETC.	- At least 2 applications during the early development period	100-150 g	400-500 g
CHICKPEAS	- Minimum of 2 applications at the beginning of flowering	100-150 g	400-500 g



EU FERTILIZERS
NK FERTILIZER SOLUTION

NATUREFERT

GUARANTEED CONTENT

	%W/W			
	18-18-18	15-30-15	5-5-40	16-8-24
Total Nitrogen (N)	18	15	5	16
Urea Nitrogen (N)	11,5	8	-	11
Ammonium Nitrogen (NH ₄ -2)	6,5	7	5	5
Water-Soluble Phosphorus Pentoxide (P ₂ O ₅)	18	30	5	8
Water-Soluble Potassium Oxide (K ₂ O)	18	15	40	24
Water-Soluble Copper (Cu)	0,03	0,02	0,03	0,02
Water-Soluble Iron (Fe)	0,05	0,05	0,05	0,05
Water-Soluble Manganese (Mn)	0,05	0,05	0,05	0,05
Water-Soluble Molybdenum (Mo)	0,001	0,001	0,001	0,001
Water-Soluble Zinc (Zn)	0,05	0,05	0,05	0,05



ORGANOMINERAL FERTILIZERS LIQUID ORGANOMINERAL FERTILIZER WITH NK

AQUA POTASOL

GUARANTEED CONTENT	%W/W
Organic Matter	20
Total Nitrogen (N)	5
Urea Nitrogen (N)	4
Organic Nitrogen (N)	1
Water-Soluble Potassium Oxide (K ₂ O)	10
Maximum Chlorine Content (Cl)	2
pH	4,1 - 6,1

RECOMMENDATIONS

- It can be applied drip or foliar during the growing season, depending on the needs of the plants.
- It is mixed with other plant fertilizing preparations and pesticides, with the exception of Bordeaux suspension and alkaline pesticides.

USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	FOLIAR APPLICATION ml/100 L of water/ha	APPLICATION TO THE SOIL L/ha
TOMATO, EGGPLANT, PEPPER, etc.	- In seedlings (2 applications), - During the period of growth	50-80 80-100	0,8-1 L 0,8-1 L
MELON, WATERMELON, CUCUMBER, etc. (PUMPKIN)	- In the period of early development - Before the first water appears - At the beginning of the sleeve development - During the fruiting period	100-120 100-120 100-120 100-120	0,8-1 L 0,8-1 L 0,8-1 L 0,8-1 L
CHERRY, PEACH, APPLE TREE, PEAR, APRICOT, PLUM, NECTARINE, Pomegranate, etc. on deciduous trees	- Immediately after sowing - When the eyes open - After flowering or when the first fruits appear - During the fruit ripening period - After picking fruits	150-250 150-250 150-250 150-250 150-250	10-15 ml per seedling 3-5 years old 15-25 ml 5-10 years old 25-40 ml per tree 10 years and older 40-50 ml per tree 40-50 ml per tree
THE VINEYARD	- Immediately after sowing - When the eyes open - Before flowering - During the corrosion period - After harvesting	150-250 150-250 100-200 100-200 100-200	5 cubes per new root 5-8 cubes per ounce of harvest 0,8-1 L from the dropper 0,8-1 L from drip irrigation 0,8-1 L from drip irrigation
OLIVES	- Immediately after sowing - Before flowering - During the summer period	200-250 200-250 200-250	In June 1-3 or 15-20 ml per tree 3-6 trees 15-25 ml 25-40 ml for 6-12 trees 40-50 ml for 12 or more trees
COTTON	- When 8-10 leaves appear (only on rows of plants) - The period before blooming and flowering - The period of the first cocoon	150-200 150-200 100-150	Per tree 1-5 years 10-15 ml 4-8 years 15-25 ml 8-15 years old 25-40 ml 40-50 ml over 15 years old
CITRUS FRUITS	- Before flowering - After flowering - June Before watering - Immediately after harvest	150-200 150-200 150-200 200-250	The dose for seeds is 100 liters of water per 2 tons of seeds, 100-150 cubes are placed and applied. Spraying is applied 100-200 ml to the decol.
POTATOES, when water is first introduced into the soil, it is given at the rate of 1 liter per 5 decaliters	- The period of the first issue - Before the first water - At the beginning of flowering - After flowering	150-200 150-200 150-200 150-200	When planting seedlings, they are dipped in 100 liters of water with a volume of 100-150 ml and planted. There is 0,8-1 L of soil per decaliter.
CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND LETTUCE	- After 15-20 days in the period of early development (2-3 ties are made)	150-200 150-200	
STRAWBERRY	- At the end of the winter period - Before and after flowering - With an interval of 15 days during harvest	100-150 100-150 100-150	0,8-1 L/ha 2 applications
CUT FLOWERS	- Rose, at the beginning of the waking period - Bud germination - With an interval of 20 days during the development period	150-200 100-150 100-150	0,8-1 L/ha 3-4 applications
GREENHOUSES	- 2 applications during development periods	150-200	Apply 0,8-1 L 2-3 times per 1000 m ² in a greenhouse by sprinkling. Pour clean water from above.



EU FERTILIZERS

EXECAL

7-0-21+(11CaO)

GUARANTEED CONTENT	%W/W
Total Nitrogen (N)	7
Nitrate Nitrogen (N)	7
Water-Soluble Potassium Oxide (K ₂ O)	21
Water-soluble Calcium Oxide (CaO)	11
Chlorine Content (Cl)	% 12,9

USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	SOIL IRRIGATION
TOMATOES, PEPPERS, WATERMELON, CARROT, STRAWBERRY, MELON	Every 15-25 days from the moment of formation of the first fruits.	2-3 L/ha
CITRUS FRUITS	Every 15-25 days from the moment of formation of the first fruits.	2-3 L/ha
APPLE, PLUM, PEACH, APRICOT, HAZELNUT	Every 15-25 days from the moment of formation of the first fruits.	2-3 L/ha
POTATOES, ONIONS, CARROT, RADISH, SUGAR BEET	2-3 applications, starting from the 4-5 leaf stage	2-3 L/ha
PEANUTS, SOYBEANS, BEANS, BEANS, etc.	2-3 applications, starting before flowering	1-2 L/ha
WHEAT, BARLEY, etc.	During the maturation period	1-2 L/ha
SWEET CORN	The plant is under construction	1-2 L/ha
OLIVES	During the formation of the first fruits, while the seeds harden	2-3 L/ha
FORAGE CROPS, CLOVER	When the plants reach 15 cm and after harvesting	2-3 L/ha
COTTON	Apple starts before grounding	1-2 L/ha
THE VINEYARD	2-3 applications After flowering	2-3 L/ha

Care should be taken when using it on chlorine-sensitive plants.

ATTENTION

- Keep away from children and food. Use a mask, gloves and glasses during fertilization.
- Never exceed the appropriate dosage and use only when necessary.
- Avoid contact with eyes, face and skin. In case of contact, rinse with plenty of water.
- Dispose of used empty packages and do not use them for other purposes.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.



ORGANOMINERAL FERTILIZERS
LIQUID MIXTURE OF NUTRIENT MICRO-PLANTS
Copper (Cu oxide), manganese (Mn oxide) and zinc (Zn oxide)

TRiFECT

GUARANTEED CONTENT	%W/W
Total Copper (Cu)	8,5
Total Manganese (Mn)	11,5
Total Zinc (Zn)	8,5

DOES NOT CONTAIN CHLORINE

ATTENTION

- Keep away from children and food. Use a mask, gloves and glasses during fertilization.
- Never exceed the appropriate dosage and use only when necessary.
- Avoid contact with eyes, face and skin. In case of contact, rinse with plenty of water.
- Dispose of used empty packages and do not use them for other purposes.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.



USE RECOMMENDATIONS

CROP	METHOD AND TIME OF APPLICATION	SOIL IRRIGATION
TOMATOES, PEPPERS, WATERMELON, CARROT, STRAWBERRY, MELON	Every 15-25 days from the moment of formation of the first fruits.	1 L/ha
CITRUS FRUITS	Every 15-25 days from the moment of formation of the first fruits.	1 L/ha
APPLE, PLUM, PEACH, APRICOT, HAZELNUT	Every 15-25 days from the moment of formation of the first fruits.	1 L/ha
POTATOES, ONIONS, CARROT, RADISH, SUGAR BEET	2-3 applications, starting from stage 4-5 leaves	1 L/ha
PEANUTS, SOYBEANS, BEANS, BEANS, etc.	2-3 applications, starting before flowering	1 - 1,5 L/ha
WHEAT, BARLEY, etc.	During the maturation period	1 - 1,5 L/ha
SWEET CORN	The plant is under construction	1 - 1,5 L/ha
OLIVES	During the formation of the first fruits, while the seeds harden	1-2 L/ha
FORAGE CROPS, CLOVER	When the plants reach 15 cm and after harvesting	1-2 L/ha
COTTON	Apple starts before grounding	1-2 L/ha
THE VINEYARD	2-3 applications After flowering	1 L/ha

ORGANIC PRODUCTS LIQUID ORGANIC FERTILIZER OF PLANT ORIGIN

FULLTECH

GUARANTEED CONTENT

	%W/W
Organic Matter	40
Organic Carbon	20
Total Nitrogen (N)	2
Water-Soluble Potassium Oxide (K ₂ O)	4
pH	4,1-6,1


USE RECOMMENDATIONS

CROP	APPLICATION PERIOD	FOLIAR APPLICATION ml/100 l of water/ha	APPLICATION TO THE SOIL l/ha
TOMATO, EGGPLANT, PEPPER, etc.	- In seedlings (2 applications), - During the period of growth	50-80 80-100	0,8-1 L 0,8-1 L
MELON, WATERMELON, CUCUMBER, etc. (PUMPKIN)	- In the period of early development - Before the first water appears - At the beginning of the sleeve development - During the fruiting period	100-120 100-120 100-120 100-120	0,8-1 L 0,8-1 L 0,8-1 L 0,8-1 L
CHERRY, PEACH, APPLE TREE, PEAR, APRICOT, PLUM, NECTARINE, POME- granate, etc. on deciduous trees	- Immediately after sowing - When the eyes open - After flowering or when the first fruits appear - During the fruit ripening period - After picking fruits	150-250 150-250 150-250 150-250 150-250	10-15 ml per seedling 3-5 years old 15-25 ml 5-10 years old 25-40 ml per tree 10 years and older 40-50 ml per tree 40-50 ml per tree
THE VINEYARD	- Immediately after sowing - When the eyes open - Before flowering - During the corrosion period - After harvesting	150-250 150-250 100-200 100-200 100-200	5 cubes per new root 5-8 cubes per ounce of harvest 0,8-1 L from the dropper 0,8-1 L from drip irrigation 0,8-1 L from drip irrigation
OLIVES	- Immediately after sowing - Before flowering - During the summer period	200-250 200-250 200-250	In June 1-3 or 15-20 ml per tree 3-6 trees 15-25 ml 25-40 ml for 6-12 trees 40-50 ml for 12 or more trees
COTTON	- When 8-10 leaves appear (only on rows of plants) - The period before blooming and flowering - The period of the first cocoon	150-200 150-200 100-150	
CITRUS FRUITS	- Before flowering - After flowering - June Before watering - Immediately after harvest	150-200 150-200 150-200 200-250	Per tree 1-5 years 10-15 ml 4-8 years 15-25 ml 8-15 years old 25-40 ml 40-50 ml over 15 years old
POTATOES, when water is first introduced into the soil, it is given at the rate of 1 liter per 5 decaliters	- The period of the first issue - Before the first water - At the beginning of flowering - After flowering	150-200 150-200 150-200 150-200	The dose for seeds is 100 liters of water per 2 tons of seeds, 100- 150 cubes are placed and applied. Spraying is applied 100-200 ml to the decol.
CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND LETTUCE	- After 15-20 days in the period of early development (2-3 ties are made)	150-200 150-200	When plantings seedlings, they are dipped in 100 liters of water with a volume of 100-150 ml and planted. There is 0,8-1 L of soil per decaliter.
STRAWBERRY	- At the end of the winter period - Before and after flowering - With an interval of 15 days during harvest	100-150 100-150 100-150	0,8-1 L/ha 2 applications
CUT FLOWERS	- Rose, at the beginning of the waking period - Bud germination - With an interval of 20 days during the development period	150-200 100-150 100-150	0,8-1 L/ha 3-4 applications
GREENHOUSES	- 2 applications during development periods	150-200	Apply 0,8-1 L 2-3 times per 1000 m ² in a greenhouse by sprinkling. Pour clean water from above.

ATTENTION

- Keep away from children and food.
- Read the label before use.
- Do not use used empty packaging for other purposes and do not destroy it.
- Use protective gloves and a mask.
- Wash your hands with plenty of water after use.
- Do not swallow. If swallowed, consult a doctor.
- Avoid contact with skin and eyes, in case of contact, rinse with plenty of water.
- It is recommended to use fertilizers after analyzing the soil and/or leaves.





NOTES:

A series of horizontal dotted lines for taking notes, spanning the width of the page below the 'NOTES:' header.



KİMYA TARIM ÜR. PAZ. SAN. VE TİC. LTD. ŞTİ.
Newagri Kimya Tarım Ür. Paz. San. ve Tic. Ltd. Şti.
Aşağıoba Mh. S.S. Toptancılar Birliği Sitesi
128 Ada B Blok No: 2 Döşemealtı / ANTALYA
Tel: 0 242 257 0 444 Fax: 0 242 221 24 00
info@newagri.com.tr • www.newagri.com.tr